White Paper Submission

Title: Constraints on Biogenesis

Author: Ian J. Miller (Dr)

Phone: 64 – 4 - 5650953

Retired.

email: ian.miller@xtra.co.nz

Introduction: Given that we have no real idea how planetesimals form, I have proposed that chemistry is the cause of initiation of planetary accretion, and the temperature dependence leads to compositional differences in bodies at different radial distance from the sun. This submission offers two suggested lines of investigation that follow from that premise, and from two published papers that in one case has been completely overlooked by the astrobiology community (through near impossibility of discovering it) and one that appears to have been forgotten. In each case the proposals depend on interpretation of what follows from the observations, but if what I expect is observed, it would make radical changes to what we think. The author is not applying for funding.

Proposal 1 Dig on Mars.

Three unresolved issues could be resolved from one premise, and if correct predictable residues should remain on Mars.

The first is that on Mars there is evidence of water flow, but there is no reasonable scenario given expected temperatures how that could occur. It is difficult to come up with a scenario where greenhouse gases might suffice that is consistent with current knowledge. While many examples of water flow on Mars might be explicable by geothermal or impact heat, some appear not. A case has been made (Parker *et al.*, 2001) that fluid flowed from the Surlus, Palaeopus and Dzigai Valles through a lake in Argire into a system forming lakes at Holden and the Ladon Valles, thence to the Margaritifer Valles, where ice was deposited. What is odd about this is the fluid started at some of the highest and coldest parts of Mars and stopped near the equator, and yet could still have flowed downhill. One possibility is that something was dissolved in the water to lower its freezing point, but was eventually lost as the water got warmer. It cannot be solid because supplies at a point would run out. There is visual evidence that at Eastern Hellas the Dao and the Harmakhis Valles change from prominent and sharp channels to diminished and muted flows that resemble terrestrial marine channels beyond river mouths, which implies there was a standing body of fluid. While the original heat would probably be volcanic, something had to lower the melting point of ice to maintain such a potentially large standing body of water. The only gas that makes sense is ammonia, which is spontaneously absorbed by ice and liquefies it down to -80 °C.

The second involves biogenesis. While there is more to it than this, for life to evolve it must start with RNA, the reason being that accurate information transfer is required, although not totally accurately, otherwise evolution is too slow. For that to happen, we need the nucleobases we have. (In principle there are further possibilities, such as orotic acid, but for the purposes here I shall ignore that because they make no difference to this overall argument.) To get those bases we need ammonium cyanide and cyanoacetylene. Again, it could be argued that there are other ways of getting to such bases, but the point I am making here is that there are no known ways of getting to such bases or alternatives in significant amounts without a reduced or partially reduced volatiles. It is almost inconceivable that we can get to life without reasonable amounts of ammonia and hydrogen cyanide being generally present. If they were, the question then is, where did they go? The amount of atmospheric nitrogen suggests that if there were sufficient for the fluid flows, the residues would be below the

surface, and may well also have chemicals on the way towards life. In principle, early precursors, as chemical fossils, may still be there.

The standard approach to the volatiles on the rocky planets is they came from carbonaceous chondrites, but there are difficulties with this. Two of the simplest ones are:

- (a) The ratios of the volatiles on the various rocky bodies are all different. Compared with Earth, Venus is slightly carbon-rich and significantly nitrogen rich, but very water poor. The high D/H ratio is argued to arise from photodissociation of water from the equivalent of Earth's oceans, or thereabouts, but there is no sign of the required oxygen. Mars is carbon poor, and extremely nitrogen poor.
- (b) There are no signs of the chondrite remains on Mars or the Moon, yet if Earth and Venus were bombarded with such chondrites, some must have struck Mars and the Moon. Neither of these have surface subduction, so such chondrites should have left remains.

Alternative Proposed Mechanism

Very briefly, the water on the rocky planets was accreted as silicate or aluminosilicate hydroxyls, and the water liberated by heat as the planet gravitationally collapsed. The nitrogen was accreted as solid nitrides or cyanides, or similar compounds, such as cyanamids formed in the heat of the accretion disk. Carbon was similarly accreted as carbides, or solid carbonaceous material. Some of the water reacted with these to form ammonia, methane, and a raft of organics that were the biochemical precursors. Venus had less water than Earth because it was hotter where it accreted. The D/H ratio on Venus is due to the chemical isotope effect during these reactions, which preferentially puts hydrogen on the volatile, or as H₂. The atmosphere would be mainly methane, and some nitrogen. Ammonia would promptly dissolve in any surface water or ice. The methane is subsequently oxidised to carbon dioxide by photochemical degradation with water providing the oxygen (and further enhancing the deuterium). The initial D/H ratio would be that of the accretion disk; the present one is determined by the concentration through the chemical isotope effect while forming volatiles, but diluted by the available water. The ammonia would dissolve in water/ice and this fluid is suggested to be what caused the fluid flow on Mars.

Support for Ammonia

The above proposal would generally be rejected because it is argued that ammonia is rapidly degraded by UV light. Thus according to Sagan and Chyba (1997) ammonia on Earth could be destroyed photolytically within decades and hence even if it were present initially, it is irrelevant. However, there is the possibility of some protection, through smogs, etc, as are found on Titan. I am unaware of data on UV photolysis of ammonia in water, but many metal salts would offer UV screening.

However, more compelling is that such rapid removal did not occur on Earth.

Seawater occluded in rock near hydrothermal vents at Barberton 3.2 Gy BP has been analysed (de Ronde *et al.* 1997), and found to contain the following gases in mmol/mol: carbon dioxide (3.3-10), nitrogen/carbon monoxide/argon/oxygen (0.4-0.9), methane (0.1-0.2), COS (0.03-0.25), C₂-C₄ hydrocarbon gases, (ca 50 ppm). Cations detected in acidified solutions (used to dissolve the rock) included ammonium at concentrations of up to 25.4 mmol/L, which is three orders of magnitude greater than current sea concentrations and approaches the level of potassium. The ammonia levels may have been greater because the salt concentrations were very high,

explicable by the geothermal heat that was the cause of trapping the seawater also boiled off some water before closure. There is no other obvious reason why seawater of the time should have a far higher salt concentration than now. If water was boiled off, some ammonia would have left too. If we assume that the nitrogen to water ratio was the same then as now, then very roughly 10% of the planet's nitrogen remained in the form of ammonia.

Ammonia would not last forever on Mars, but given that any specific Martian fluid flows seldom lasted more than 100,000 years, ammonia provides a reason why. In this context, the oxidation of the methane produces carbon dioxide, and this reacts with ammonia to form ammonium carbonate, which in turn may dehydrate to urea. If ammonia was what stabilized the water as a fluid, such urea would eventually settle out as a solid, and either is still there, in some form or other (chemistry is not very good at predicting what slow reactions could occur over billions of years) or it has volatilised, but that is unlikely because of the serious nitrogen deficiency. Ammonia will also concentrate certain transition metal oxides as ammines, but this is less definitive because acid does the same thing, as at Meridiani. Thus if there were nitrogen there in that form in sufficient volumes to fluidize that much water, it should still be there. If it is not, then Mars was always nitrogen deficient.

Accordingly, the proposal is that a site be found where water could reasonably have lain, and the dust level of overburden does not exceed the plausible depth that a rover could dig.

Option 2: Chemical laboratory work to address, Why ribose?

Vesicles have been made from fatty acids and glycerol esters that acid catalysed condensation by sequential wetting and drying (Milshteyn *et al.* 2018). These vesicles were stable even at pH 3.3, and were stable to the drying, even on rocks, however they were not stable to seawater. Nucleotides could be condensed to polymers of between 10 – 100 mers (Ross and Deamer, 2016). This condensation reaction is endoergic by about 13.8 kJ/mol, but the vesicle wet/dry cycling provides stabilizing factors of 20 – 40 kJ/mol. It has been shown (Hassenkam *et al.* 2020) that sequential wet/dry cycles of vesicles containing AMP and UMP lead to polymer strands, which led to the conclusion that RNA-like rings and structures with sizes sufficient to fold into ribozymes could have been formed abiogenically. Thus we can see how AMP, UMP, GMP and CMP could lead to the origin of life. The problem then is, how to get to AMP, UMP, GMP and CMP? This, in turn, reduces to, how to make phosphate esters in water in a "one pot" type reaction without significant changes of conditions. Phosphate will esterify alcohols at about 200 °C, but the esters hydrolyse at that temperature in the presence of excess water, and saccharides caramelize or decompose well before that. Further, it should be done in a vesicle so the rest of the RNA synthesis can take place.

One proposed route (Ponnamperuma *et al.*, 1963) proceeded through purines absorbing UV radiation at about 250 nm wavelength in the presence of ribose and a phosphate. Thus adenine gave AMP and ATP. There are three major problems with the Ponnamperuma mechanism. It is highly unlikely this UV window would be available at suitable wavelengths, especially those that would permit uracil to absorb, because upper atmosphere photochemistry would screen it. The second is that it has been shown (Shuttlefield *et al.*, 2011) to be highly desirable that this radiation is *not available* because UV radiation at 250 nm striking oxide semiconductors such as titanium dioxide, and probably ilmenite and haematite, oxidizes chloride to

perchlorate. This type of reaction is quite likely to be the source of the perchlorate on Mars, and it is clear that such strong oxidations would not be conducive to the emergence of life, particularly since iron and titanium oxides are not rare chemicals. The third problem is the solubility of phosphate. This will generally appear as something like calcium hydroxylapatite, which is insoluble in water. However, the presence of urea and ammonium formate increases the solubility, and also apparently enhances the chances of phosphorylation (Burcar *et al.* 2016), particularly in the presence of magnesium sulphate. Ammonium formate is a hydrolysis product of hydrogen cyanide. Also, it has been found that cyanide itself, in the presence of urea or as an iron cyanide complex also improves phosphate solubility, while phosphate may well have been present in more mobile phosphates, such as magnesium ammonium phosphate (Burcar *et al.* 2019), and these slowly phosphorylated adenosine. It was unclear whether photochemistry had any involvement here but the samples would have been in the light, and cyanide complexes with ferrous could have been light absorbers.

RNA is not the only plausible nucleic acid and a number of pentose pyranoside nucleic acids, including ribopyranoside, form duplexes with stronger binding energy than RNA (Eschenmoser, 1999), so why did nature select ribose, which is made in poor yield in most abiogenic syntheses? (The complexation of aldose sugars with silicate strongly favours ribose by an order of magnitude, and equally, hexoses such as glucose, mannose and galactose do not form such complexes (Lambert *et al.* 2004, 2010). This would favor the origin of life forming around fumaroles.) The proposed answer is because ribose is unusual as a sugar in that in aqueous solution it has about 18% in the furanose form (Cocinero *et al.* 2012), while other sugars show very little furanose form. But why ribofuranoside, when there is more pyranoside that can participate in duplexes?

How do the phosphate esters form, and why at C-5? In the Ponnampereuma route, it is not directly photochemical as the excited state cannot be transmitted through σ bonds, but mechanical energy can, and the furanose form alone has the flexibility to do this. The simplest answer (but not necessarily the correct one) is the furanoside flexes and sufficient mechanical energy is focused at O5 to provide the vibrational motion equivalent to a hydroxyl at 200 °C. The water does not hydrolyse the product because it is at room temperature. If so, a mix of sugars in the Ponnampereuma reaction will only yield primary ribofuranoside phosphate ester formation. However, the UV required is too harsh. The alternative is that if the reaction is carried out in a vesicle, with porphyrin or some similar dye, the mechanical energy can be transmitted through ribose from radiationless decay of the excited states. The proposal is, first repeat the Ponnampereuma reaction then with ribose plus other sugars. If the proposal is correct, the other sugars should not participate in the primary reaction. If so, and if mechanical energy can be transmitted from the vesicle walls, the vesicles with a porphyrin or suitable dye should permit the formation of nucleotides with visible radiation. If so, coupled with the geologic requirements to form ribose, there should be very clear limits to where life can form, and furthermore, the very initial life form would have the characteristics that would eventually lead to photosynthesis as we know it.

References:

- Burcar, B. and 6 others. 2016. Darwin's Warm Little Pond: A One-Pot Reaction for Prebiotic Phosphorylation and the Mobilization of Phosphate from Minerals in a Urea-Based Solvent. *Angew. Chem. Int. Ed.* 55: 13249 – 13253.

- Burcar, B. and 7 others. 2019. A Stark Contrast to Modern Earth: Phosphate Mineral Transformation and Nucleoside Phosphorylation in an Iron- and Cyanide-Rich Early Earth Scenario. *Angew. Chem. Int Ed.* 58: 16981 – 16987.
- Cocinero, E. J. and 6 others. 2012. Ribose found in the gas phase. *Angew. Chemie Internat. Ed.* 51: 3119-3124.
- de Ronde, C., Channer, D. M. de R., Faure, K., Bray, C. J. and Spooner, T. C., 1997. Fluid chemistry of the Archaean seafloor hydrothermal vents: implications for the composition of circa 3.2 Ga seawater. *Geochim. Cosmochim. Acta* 61: 4025-4042.
- Eschenmoser, A. 1999. Chemical etiology of nucleic acid structure. *Science* 284: 2118 – 2124.
- Hassenkam, T., Damer, B., Mednick, G. Deamer, D. 2020. Viroid-sized rings self-assemble from mononucleotides through wet-dry cycling: implications for the origin of life. <https://www.biorxiv.org/content/10.1101/2020.04.28.064972v1>
- Lambert, J. A., Gurusamy-Thangavelu, S. A., Ma, K. 2010. The silicate-mediated formose reaction: bottom-up synthesis of sugar silicates. *Science* 327: 984-986.
- Lambert, J. A., Lu, G., Singer, S. R., Kolb, V.M. 2004. Silicate Complexes of Sugars in Aqueous Solution. *J. Amer. Chem. Soc.* 126: 9611 – 9625.
- Milshteyn, D., Damer, B., Havig, J., Deamer, D. 2018. Amphiphilic compounds assemble into membranous vesicles in hydrothermal hot spring water but not in seawater. *Life* 8: (2) 11. <https://doi.org/10.3390/life8020011>
- Parker, T. J., Clifford, S. M., Bannert, W. B. 2001. Argyre Planitia and the Mars global hydrological cycle. *Lunar Planet. Sci. Conf. XXXI*, Abstr. 2033.
- Ponnamperuma, C., Sagan, C., Mariner, R., 1963. Synthesis of adenosine triphosphate under possible primitive earth conditions. *Nature* 199: 222-226.
- Ross, D. S., Deamer, D. 2016. Dry/Wet Cycling and the Thermodynamics and Kinetics of Prebiotic Polymer Synthesis. *Life* 6: 28 (12 pp) doi:10.3390/life6030028.
- Sagan, C. and Chyba, C., 1997. The early faint sun paradox: organic shielding of ultraviolet-labile greenhouse gases. *Science* 276: 1217-1221.
- Shuttlefield, J. D., Sambur, J. B., Gelwicks, M., Eggleston, C. M., Parkinson, B. A. 2011. Photooxidation of chloride by oxide minerals: implications for perchlorate on Mars. *J. Am Chem. Soc.* 133: 17521-17523.